SESAPS 2007 – Nashville, TN

Proposed operator description of four dimensional Euclidean quantum gravity

(SES07-2007-000015)

Author: Jens Koeplinger
Date: 09 November 2007
Contact: jens@prisage.com
Publications and preprints: http://www.jenskoeplinger.com/P

(CC/by 3.0) 2007 (http://creativecommons.org/licenses/by/3.0/)

Context: Quantum gravity

- "4D Euclidean quantum gravity" (EQG)
  - Gibbons, Hawking: "Euclidean Quantum Gravity" [1]
    - ed., (1993), Singapore: World Scientific
  - Field theory / second quantization
  - Expectation: Operator description of 4D EQG unlikely
  - Suggestion for realization [2]: anti-de Sitter (AdS) metric
- Research problem of this talk: Find operator (first quantization) description

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Context: Standard Model

Operator description (1^{st} quantization)	Field theory (2^{nd} quantization)
	Strong force Weak force
Electromagnetism	Electromagnetism

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Proposed: Standard Model + EQG

Operator description (1^{st} quantization)		Field theory (2^{nd} quantization)
		Strong force Weak force
		↑
Electromagnetism	↔	Electromagnetism
↓		↓
Proposed here: 4D EQG	↔	4D EQG

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Computation concepts

Operator description
(1st quantization)

Field theory
(2nd quantization)



(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Motivation

- Why normed algebra?
  - Use metric to define relativity: Different frames of reference are equivalent when energy-momentum vector norm is invariant.
- Why non-associative algebra?
  - Consistent with known physics: Octonionic expression of QED (Gogberashvili [3,4], Köplinger [8])
  - Octonionic Proca-Maxwell equations [9]
  - "New" (non-matrix) symmetry transformations possible (Köplinger [5-8])

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Cayley-Dickson constructs ...

(real)	$1\ i_1\ i_2\ i_3\ i_4\ i_5\ i_6\ i_7$	
Complex	■ ■	
Quaternion	■ ■ ■ ■	
Octonion	■ ■ ■ ■ ■ ■ ■ ■	

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

... and "complexified" algebras

(real)	$1\ i_1\ i_2\ i_3\ i_4\ i_5\ i_6\ i_7\ i_0\varepsilon_1\varepsilon_2\varepsilon_3\varepsilon_4\varepsilon_5\varepsilon_6\varepsilon_7$	(complex)
Complex	■ ■ ■ ■	Tessarine (or conic quat., bicomplex num)
Quaternion	■ ■ ■ ■ ■ ■ ■ ■	Biquaternion (or compl. quat., conic octonion)
Octonion	■ ■ ■ ■ ■ ■ ■ ■ ■ ■ ■ ■ ■ ■ ■ ■	Complex octonion (or conic sedenion)

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Spin ½ particle w/o field

	$1\ i_1\ i_2\ i_3\ i_4\ i_5\ i_6\ i_7\ i_0\varepsilon_1\varepsilon_2\varepsilon_3\varepsilon_4\varepsilon_5\varepsilon_6\varepsilon_7$	
Gogberashvili [3]	■ ■ ■ ■ ■ ■ ■ ■	Dirac eqn
Köplinger [5]	■ ■ ■ ■ ■ ■ ■ ■	Dirac eqn

$$\Psi_{\text{hyp8}} := \left(\psi_0^{\text{r}}, \psi_0^{\text{i}}, \psi_1^{\text{r}}, \psi_1^{\text{i}}, \quad \psi_2^{\text{r}}, -\psi_2^{\text{i}}, -\psi_3^{\text{r}}, -\psi_3^{\text{i}}\right)$$

$$\nabla_{\text{hyp8}} := (-m, \partial_0, 0, 0, \quad 0, -\partial_3, \partial_2, -\partial_1)$$

$$\nabla_{\text{hyp8}}\, \Psi_{\text{hyp8}} = 0$$

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Spin ½ particle with field

	$1\ i_1\ i_2\ i_3\ i_4\ i_5\ i_6\ i_7\ i_0\varepsilon_1\varepsilon_2\varepsilon_3\varepsilon_4\varepsilon_5\varepsilon_6\varepsilon_7$	
Gogberashvili [3]	■■■■ ■■■■	Dirac eqn
Gogberashvili [4]	■■■■ ■■■■	EM
Köplinger [5]	■■■■ ■■■■	Dirac eqn
Köplinger [6]	"New" (non-matrix) symmetry trans.: ■■■■■■■■	"hvy"Dirac
Köplinger [7]	Define consistent "relativity": ■■■■■■■■■■■■■■■■	GRT equiv
Köplinger [8]	Add EM field consistent with QED: ■■■■■■■■■■■■■■■■	EM & GR

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Consistency (non-quantum)

- How would we observe a complex metric that is both Euclidean and Minkowskian?
  - Large body, non-quantum effects don't mix forces; we can first look at Euclidean and Minkowski metrics separately, and then project the effects from Euclidean geometry onto Minkowskian geometry.
  - Supporting argument: Our daily experience is dominated by Electromagnetism and Minkowski space-time, other forces may appear as a modification thereof.

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Model: NatAliE equations

- "Naturally aligned elementary equations"
  - Treat gravity on global 4D Euclidean space-time, separately from all other forces.
  - For this space-time only, develop relativity in direct correspondence to Special Relativity (SRT):
    - Unaccelerated frames of reference are equivalent.
    - "Heavy" Lorentz transformation conserves 4D Euclidean length element and energy-momentum.
    - As little changes as possible compared to SRT.
  - In the SRT equation of motion, modify the gravitational field geometry accordingly.

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Result (non-quantum)

- NatAliE equations become linearized field equations from General Relativity (GRT) [7]
  - Since we did not look at field self-interaction, this is the desired result (for consistency).
  - Known "bootstrap" method yields covariant GRT.
  - Conclusion: NatAliE is consistent with experimental results for large body, non-quantum physics [7].
- Now: generalize for quantum physics

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Model for quantum gravity

- Perform operator transition [8]:
  - Conic sedenions (complex octonions) as enveloping algebra that contains both Euclidean and Minkowskian geometry.
  - Add EM field consistent with QED.
  - Introduce "mixing angle" to transition between electromagnetism and gravity.
- Relativity warranted by invariant modulus (norm) of the energy-momentum vector.

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Spin ½ particle with field

	$1\ i_1\ i_2\ i_3\ i_4\ i_5\ i_6\ i_7\ i_0 \varepsilon_1 \varepsilon_2 \varepsilon_3 \varepsilon_4 \varepsilon_5 \varepsilon_6 \varepsilon_7$	
Köplinger [8]	■ ■ ■ ■ ■ ■ ■ ■ ■ ■ ■ ■ ■ ■ ■ ■	EM & GR

$$
\begin{aligned}
\nabla_{Q1} &:= (0, \partial_0, 0, 0, 0, 0, 0, 0, \quad 0, eA_0, 0, 0, 0, 0, 0, 0) \\
\nabla_{Q2} &:= (0, 0, 0, 0, 0, \partial_3, -\partial_2, \partial_1, \quad 0, 0, 0, 0, 0, eA_3, -eA_2, eA_1) \\
\nabla &:= \nabla_{Q1} + \exp(i_0 \alpha) \nabla_{Q2}
\end{aligned}
$$

$$
\begin{aligned}
\Psi_{Q1} &:= (\psi_0^{\mathrm{r}}, \psi_0^{\mathrm{i}}, \psi_1^{\mathrm{r}}, \psi_1^{\mathrm{i}}, 0, 0, 0, 0, \quad 0, 0, 0, 0, 0, 0, 0, 0) \\
\Psi_{Q2} &:= (0, 0, 0, 0, -\psi_2^{\mathrm{r}}, \psi_2^{\mathrm{i}}, \psi_3^{\mathrm{r}}, \psi_3^{\mathrm{i}}, \quad 0, 0, 0, 0, 0, 0, 0, 0) \\
\Psi &:= \Psi_{Q1} + \exp(i_0 \alpha) \Psi_{Q2}
\end{aligned}
$$

$$
(\nabla - m) \Psi = 0
$$

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Predictions

- Challenge: Quantum gravity's weakness
  - Conic sedenion formulation reduces to tensor algebra in excellent approximation for current experiments [8]. This is a challenge in validating any theory on quantum gravity.
- Initial findings
  - Backscattering is increased for elastic gravitational spin ½ scattering
  - Scattering cross section does not vanish for E --> ∞

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Outlook

- To examine:
  - Measurable effects that distinguish this model from other models on quantum gravity?
  - Vacuum effects?
  - Extendability of Gogberashvili's approach?
  - Effects at Big Bang energy levels?
  - Massive exchange particle(s)?
  - Other forces?

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Acknowledgments

My special thanks and gratitude to

Kevin Carmody,
John Shuster,
Armahedi Mahzar, and
Marek Čtrnáct

for helping me understand and utilize algebraic concepts, and for vital and fruitful discussions.

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

Appendix: Conic sedenions

Quick notes about conic sedenions (complex octonions) [10,11]

Modulus (norm): $$|z| = |a + \sum b_n i_n + \sum c_n \varepsilon_n + d| := \sqrt[4]{(a^2 + b_n^2 - c_n^2 - d^2)^2 + 4(ad - b_n c_n)^2}$$

Multiplication table:

X	1	i_1	i_2	i_3	i_4	i_5	i_6	i_7	i_0	ε_1	ε_2	ε_3	ε_4	ε_5	ε_6	ε_7
1	1	i_1	i_2	i_3	i_4	i_5	i_6	i_7	i_0	ε_1	ε_2	ε_3	ε_4	ε_5	ε_6	ε_7
i_1	i_1	-1	i_3	$-i_2$	i_5	$-i_4$	$-i_7$	i_6	$-\varepsilon_1$	i_0	ε_3	$-\varepsilon_2$	ε_5	$-\varepsilon_4$	$-\varepsilon_7$	ε_6
i_2	i_2	$-i_3$	-1	i_1	i_6	i_7	$-i_4$	$-i_5$	$-\varepsilon_2$	$-\varepsilon_3$	i_0	ε_1	ε_6	ε_7	$-\varepsilon_4$	$-\varepsilon_5$
i_3	i_3	i_2	$-i_1$	-1	i_7	$-i_6$	i_5	$-i_4$	$-\varepsilon_3$	ε_2	$-\varepsilon_1$	i_0	ε_7	$-\varepsilon_6$	ε_5	$-\varepsilon_4$
i_4	i_4	$-i_5$	$-i_6$	$-i_7$	-1	i_1	i_2	i_3	$-\varepsilon_4$	$-\varepsilon_5$	$-\varepsilon_6$	$-\varepsilon_7$	i_0	ε_1	ε_2	ε_3
i_5	i_5	i_4	$-i_7$	i_6	$-i_1$	-1	$-i_3$	i_2	$-\varepsilon_5$	ε_4	$-\varepsilon_7$	ε_6	$-\varepsilon_1$	i_0	$-\varepsilon_3$	ε_2
i_6	i_6	i_7	i_4	$-i_5$	$-i_2$	i_3	-1	$-i_1$	$-\varepsilon_6$	ε_7	ε_4	$-\varepsilon_5$	$-\varepsilon_2$	ε_3	i_0	$-\varepsilon_1$
i_7	i_7	$-i_6$	i_5	i_4	$-i_3$	$-i_2$	i_1	-1	$-\varepsilon_7$	$-\varepsilon_6$	ε_5	ε_4	$-\varepsilon_3$	$-\varepsilon_2$	ε_1	i_0
i_0	i_0	$-\varepsilon_1$	$-\varepsilon_2$	$-\varepsilon_3$	$-\varepsilon_4$	$-\varepsilon_5$	$-\varepsilon_6$	$-\varepsilon_7$	-1	i_1	i_2	i_3	i_4	i_5	i_6	i_7
ε_1	ε_1	i_0	ε_3	$-\varepsilon_2$	ε_5	$-\varepsilon_4$	$-\varepsilon_7$	ε_6	i_1	1	$-i_3$	i_2	$-i_5$	i_4	i_7	$-i_6$
ε_2	ε_2	$-\varepsilon_3$	i_0	ε_1	ε_6	ε_7	$-\varepsilon_4$	$-\varepsilon_5$	i_2	i_3	1	$-i_1$	$-i_6$	$-i_7$	i_4	i_5
ε_3	ε_3	ε_2	$-\varepsilon_1$	i_0	ε_7	$-\varepsilon_6$	ε_5	$-\varepsilon_4$	i_3	$-i_2$	i_1	1	$-i_7$	i_6	$-i_5$	i_4
ε_4	ε_4	$-\varepsilon_5$	$-\varepsilon_6$	$-\varepsilon_7$	i_0	ε_1	ε_2	ε_3	i_4	i_5	i_6	i_7	1	$-i_1$	$-i_2$	$-i_3$
ε_5	ε_5	ε_4	$-\varepsilon_7$	ε_6	$-\varepsilon_1$	i_0	$-\varepsilon_3$	ε_2	i_5	$-i_4$	i_7	$-i_6$	i_1	1	i_3	$-i_2$
ε_6	ε_6	ε_7	ε_4	$-\varepsilon_5$	$-\varepsilon_2$	ε_3	i_0	$-\varepsilon_1$	i_6	$-i_7$	$-i_4$	i_5	i_2	$-i_3$	1	i_1
ε_7	ε_7	$-\varepsilon_6$	ε_5	ε_4	$-\varepsilon_3$	$-\varepsilon_2$	ε_1	i_0	i_7	i_6	$-i_5$	$-i_4$	i_3	i_2	$-i_1$	1

Other relations:

$$e^{\varepsilon_n \alpha} = \cosh\ \alpha + \varepsilon_n(\sinh\ \alpha)$$

$$\ln \varepsilon_n = \frac{\pi}{2}(i_0 - i_n)$$

$$\varepsilon_n^{\alpha} = \frac{1}{2}[(1 - \varepsilon_n) + (1 + \varepsilon_n)e^{-\pi i_n \alpha}]$$

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com

References

[1] G. W. Gibbons and S. W. Hawking (ed), *Euclidean Quantum Gravity* (World Scientific, Singapore, 2004).

[2] S. W. Hawking, *Information loss in black holes*, Phys. Rev. D **72**, 084013 (2005).

[3] M. Gogberashvili, *Octonionic version of Dirac equations*, Int. J. Mod. Phys. A **21**, 3513-3524 (2006).

[4] M. Gogberashvili, *Octonionic electrodynamics*, J. Phys. A – Math. Gen. **39** 7099-7104 (2006).

[5] J. Köplinger, *Dirac equation on hyperbolic octonions*, Appl. Math. Comput. **182**, 443-446 (2006).

[6] J. Köplinger, *Signature of gravity in conic sedenions*, Appl. Math. Comput. **188**, 942-947 (2007).

[7] J. Köplinger, *Hypernumbers and relativity*, Appl. Math. Comput. **188**, 954-969 (2007).

[8] J. Köplinger, *Gravity and electromagnetism on conic sedenions*, Appl. Math. Comput. **188**, 948-953 (2007).

[9] N. Candemir, M. Tanişli, K. Özsaş, and S. Demir, *Hyperbolic octonionic Proca-Maxwell equations*, Z. Naturforsch. **63a**, 15-18 (2008).

[10] K. Carmody, *Circular and hyperbolic quaternions, octonions, and sedenions*, Appl. Math. Comput. **28**, 47-72 (1988).

[11] K. Carmody, *Circular and hyperbolic quaternions, octonions, and sedenions — further results*, Appl. Math. Comput. **84**, 27-48 (1997).

(CC/by 3.0) 2007 Jens Koeplinger - http://www.jenskoeplinger.com